

**Research Article****Open Access**

# Post-operative Hypothermia in Surgical Patients at University of Gondar Hospital, Ethiopia

Tadesse Belayneh<sup>1\*</sup>, Abebaw Gebeyehu<sup>2</sup> and Zewditu Abdissa<sup>3</sup><sup>1</sup>Department of Medical Anesthesiology, School of Medicine, University of Gondar, Ethiopia<sup>2</sup>Department of Reproductive Health, Institute of Public Health, University of Gondar, Ethiopia<sup>3</sup>Department of Medical Anesthesiology, School of Medicine, University of Gondar, Ethiopia**Abstract**

**Background:** Hypothermia (i.e., temperature  $<36^{\circ}\text{C}$ ) is a common incident in the immediate postoperative period. However, it is usually diagnosed and treated infrequently. Maintaining normal body temperature will reduce hospital stay, surgical site infection, postoperative blood transfusions, pressure ulcers, subjective discomfort and mortality. The aim of this study was to identify magnitude and factors associated with postoperative hypothermia.

**Methods:** Hospital based cross sectional study was conducted at University of Gondar Hospital. All consecutive postoperative (N= 384) patients that underwent scheduled or emergency surgery were included. Tympanic membrane temperature was measured before surgery and on arrival at recovery room via clinical nurses. Analysis was done using version 20 SPSS and binary logistic regression was utilized to identify associated factors.

**Result:** The incidence of postoperative hypothermia at Recovery Room was 30.72%. Of these, 30 (27.96%) and 43 (34.44%) were mild and moderate respectively, with no severe hypothermia. Surgical patient operated under general anesthesia was more likely to develop postoperative hypothermia than spinal anesthesia (AOR, 1.84; 95% CI, 1.17; 3.24). Additionally, those with higher ASA physical status were at higher risk of postoperative hypothermia (AOR, 1.99; 95% CI, 1.16; 3.44). Temperature was not routinely taken by health care providers in the preoperative and intraoperative period.

**Conclusion and recommendation:** Incidence of postoperative hypothermia at Recovery Room was high. Patient with high ASA physical status and operated under general anesthesia were significant factors for the development of postoperative hypothermia. Anesthesia providers along with perioperative surgical team should participate to reduce the occurrence of postoperative hypothermia with early prediction and diagnosis, especially in high risk groups.

**Keywords:** Postoperative hypothermia; Recovery room; Surgical patients

**Introduction**

Hypothermia is defined as a body temperature of  $<36^{\circ}\text{C}$  ( $96.8^{\circ}\text{F}$ ) and may be classified as mild ( $35.0^{\circ}\text{C}$ - $35.9^{\circ}\text{C}$ ), moderate ( $34.0^{\circ}\text{C}$ - $34.9^{\circ}\text{C}$ ) and severe ( $\leq 33.9^{\circ}\text{C}$ ) [1-3]. Despite, overlooked as an inevitable consequence, it is usually a typical incident of surgical patients [4,5]. It is a frequent occurrence that affects more than 70% of patients undergoing surgery and anesthesia. Moreover, it is the most cause of recovery room admission in the immediate postoperative period [6]. A body temperature change of  $1\text{-}3^{\circ}\text{C}$  leads multiple physiological derangements [7,8], such as, change in protein metabolism [9] that compromise healing and platelet dysfunction [10] which leads to bleeding and postoperative transfusions [11], arrhythmia [12], prolonged hospital stay [13], surgical site infection and pressure ulcers [14,15]. It also lowers patient satisfaction with the services [16,17] and causes subjective discomfort due to cold and shivering [18].

In the postoperative period, patients might be hypothermic due to different reasons. This may cause prolonged hospital stay, surgical site infection, postoperative blood transfusions, pressure ulcers, subjective discomfort and mortality [13-16]. The occurrence of postoperative hypothermia is associated with older age [16], female, emergency surgery [19], higher American Society of Anesthesiology physical status, major surgical procedure, amount of intravenous or blood replaced, longer duration of anesthesia or surgery, operating room temperature [20,21], preoperative body temperature and anesthetic technique [22].

Temperature monitoring provides an early detection of postoperative hypothermia. As Dubois described, there are two (core and periphery) thermal compartments for temperatures of the human body [23]. The pulmonary artery, nasopharyngeal, esophageal and tympanic temperatures measurements are the best temperature monitoring sites [24], but temperature from bladder, axilla, skin and rectal are slowest to change and underestimate the magnitude of alteration in body temperature [25,26]. Although ear thermometer is easy and fast, its accuracy has always been controversial. It has been noticed that an ear thermometer reading varies from  $1$  to  $2^{\circ}\text{C}$  below and above the temperature than rectal and oral thermometers [27]. Conversely, tympanic temperature measurements seem to be the best alternative to rectal and infrared skin measurements with appropriate use. The reading can be affected by place of insertion, size of ears (intended for people older than 1 year of age), fluid and exudate [28]. The device was designed for patients suffering from accidental hypothermia [29].

**\*Corresponding author:** Tadesse Belayneh, Department of Medical Anesthesiology, School of Medicine, University of Gondar, Ethiopia, Tel: 251-918777496; E-mail: [tadbel20@gmail.com](mailto:tadbel20@gmail.com)

**Received** August 14, 2014; **Accepted** October 29, 2014; **Published** November 20, 2014

**Citation:** Belayneh T, Gebeyehu A, Abdissa Z (2014) Post-operative Hypothermia in Surgical Patients at University of Gondar Hospital, Ethiopia. J Anesth Clin Res 5: 461. doi: [10.4172/2155-6148.1000461](https://doi.org/10.4172/2155-6148.1000461)

**Copyright:** © 2014 Belayneh T, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

In our hospital, surgical patients are transferred to recovery room almost naked with a simple cover of surgical drapes and gowns postoperatively. This will expose the patient to the cold environment that causes heat loss and adverse effects. Anesthesia providers along with postoperative surgical team remain diligent in implementing a variety of interventions to minimize postoperative hypothermia. In an effort to decrease the occurrence of postoperative hypothermia, it would be helpful to identify and correct significant associated factors. Therefore, the aim this study was to assess magnitude and associated factors of postoperative hypothermia at Recovery Room of University of Gondar Hospital, Northwest Ethiopia.

## Methods

Hospital based cross sectional study was conducted from January to March 2014 at University of Gondar Hospital. According to review of surgical registry book, 14082 patients were operated in 2012/13. Postoperatively, patients are admitted in the Recovery Room until their vital signs are stabilized before transferred to wards.

All consecutive patient that underwent scheduled or emergency non-cardiac surgery were included while patient with fever, sepsis and procedure performed under local anaesthesia were excluded from the study. A single population proportion formula was used to estimate the sample size with the following assumptions: P as 50%, considering 95% confidence interval and 5% margin of error ( $d=0.05$ ). The total sample size then, was 384 patients.

Individual tympanic membrane temperature was measured before surgery and on arrival at RR. It was measured using reusable infrared thermometer (Thermos Scan<sup>®</sup> Type 6014 Pro 3000 WelchAllyn with an accuracy of  $\pm 0.03^{\circ}\text{C}$  in the range of  $20^{\circ}\text{C}$ - $42.2^{\circ}\text{C}$ ). This thermometer is considered ideal because the tympanic membrane and hypothalamus share an arterial blood supply originating from the carotid artery; therefore, the tympanic membrane is considered to directly reflect core temperature [30]. All patients were examined to exclude ear infection and occluding ear wax was cleared. The probe of the thermometer was inserted into the external auditory meatus (either the left or right ear) by pulling the pinna backward, and directing the probe towards the eye. The instrument was calibrated in accordance with the manufacturer's manual of instruction before each reading. The ambient OR temperature was also measured using the thermocouple room thermometer. Postoperatively patients were classified as hypothermic ( $<36^{\circ}\text{C}$ ) or normothermic ( $\geq 36^{\circ}\text{C}$ ) [1,31]. All other data were collected from patient medical record.

All responses to the questionnaires were coded, entered and analyzed using SPSS Version 20. Bivariate logistic regression was employed to determine associated factors between variables. P-value and 95% confidence interval was used to check statistical significance. The distribution normality of each variable was analyzed in the Kolmogorov-Smirnov test.

Ethical clearance was taken from Gondar College of Medicine and Health Sciences Research Ethical Review committee. Official permission was also obtained from University of Gondar Hospital. Preoperatively, participants were told individually about the purpose of the study and verbal informed consent was taken. The right of participants to withdraw from the study at any time, without any pre-condition was kept, and disclosed to respondents. Confidentiality of the information was assured. All hypothermic patients were managed with external rewarming measures that include more synthetic blanket covers and electrical heater.

## Results

### Sociodemographic characteristics of postoperative hypothermia

A total of 384 patients with age range from 1 to 83 years (median, 28.0) were included in the study. Among these, 52.1% were females and 68.5% operated with general anesthesia. The median Operation Room ambient temperature was  $20.0^{\circ}\text{C}$  with a range of  $8.8^{\circ}\text{C}$ . Duration of surgery was ranged from 30-210 minutes (median, 85 minutes); (Table 1).

### Magnitude of postoperative hypothermia

The incidence of postoperative hypothermia immediately at recovery room was 30.72%. Of these, 30 (27.96%) and 43 (34.44%) were mild and moderate respectively with no severe hypothermia noticed. The median (Range) temperature was  $36.80$  ( $0.96$ ) $^{\circ}\text{C}$  in preoperative patient but  $36.10$  ( $0.83$ ) $^{\circ}\text{C}$  in the postoperative cases. The paired t-test showed significant difference in pre and post operative temperature measurements. Patients with hypothermia were more frequently in age of 25-50 years (52.5%), male gender (50.8%), with ASA I (79.7%), having a normal preoperative temperature (81.4%) and underwent schedule operation (62.7%). Temperature was not monitored preoperatively and intraoperatively in the operation room by nurses and anesthesia providers respectively. All crystalloid fluids were administered without warming but blood and abdominal irrigation fluids were utilized after warming, though the temperature was not adjusted in the recommended range of body temperature. Occurrence of hypothermia was prevented using passive warming measures such as cotton sheets, drapes, gowns or blankets covering in the intraoperative period and during patient transfer to recovery room.

Characteristics	Frequency (%)
Age (years)	<25 107 (27.9%)
	25-50 205 (53.4%)
	>50 years 72 (18.8%)
ASA physical status	I 280 (72.9%)
	II 104 (27.1%)
Sex	Male 184 (47.9%)
	Female 200 (52.1%)
Preoperative temperature ( $^{\circ}\text{C}$ )	$<36.0^{\circ}\text{C}$ 76 (19.8%)
	$\geq 36.0^{\circ}\text{C}$ 308 (80.2%)
Magnitude of procedure	Moderate 84 (28.1%)
	Major 276 (71.9%)
Type of anesthesia	General 263 (68.5%)
	Spinal 121 (31.5%)
Condition of surgery	Elective 214 (55.7%)
	Emergency 170 (44.3%)
Amount of crystalloid fluids replaced intraoperatively (litre)	
	$\leq 2$ 303 (78.9%)
	$>2$ 81 (21.1%)
Duration of Anesthesia (min) (Range 35-230 min)	
	$\leq 1$ hr 57 (14.8%)
	$>1$ hr 327 (85.2%)
Duration of Surgery (min) (Range 30-210 min)	
	$\leq 1$ hr 121 (31.5%)
	$>1$ hr 263 (68.5%)
Operation room temperature	$\leq 23^{\circ}\text{C}$ 309 (80.5)
	$>23^{\circ}\text{C}$ 75 (19.5)

**Table 1:** Characteristics of patients by postoperative temperature status in Gondar town, Northwest Ethiopia, 2014 (n=384).

Characteristics		Status of temperature		Crude Odds Ratio (95% CI)	Adjusted Odds Ratio (95% CI)	P- value
		Hypothermic	Euthermic			
ASA physical status	I	94 (79.7%)	186 (69.9%)	1	1	0.01
	II	24 (20.3%)	80 (30.1%)	2.5 (1.5-4.0)	1.99 (1.16-3.44)	
Type of anesthesia	General	91 (77.1%)	172 (64.7%)	1.9 (1.3-3.1)	1.84 (1.17-3.24)	0.02
	Spinal	27 (22.9%)	94 (35.3%)	1	1	
Condition of surgery	Elective	74 (62.7%)	140 (52.6%)	1	1	0.03
	Emergency	44 (37.3%)	126 (47.4%)	0.7 (0.5-1.1)	0.56 (0.33-0.94)	

1: Reference group

**Table 2:** Factors associated with postoperative hypothermia in surgical patient of University of Gondar Hospital, North West Ethiopia, 2014 (n=384).

## Factors associated with postoperative hypothermia

The multivariate logistic regression in Table 2 indicated that patients operated under general anesthesia were 1.8 times more likely develop postoperative hypothermia than operated under spinal anesthesia (AOR, 1.84; 95% CI, 1.17; 3.24). Among the preoperative factors, emergency surgery were significant protective factors against hypothermia (AOR, 0.52; 95% CI, 0.31; 0.84). Those patient with higher ASA physical status were at higher risk of postoperative hypothermia (AOR, 1.99; 95% CI, 1.16; 3.44) than lower ASA status. No statistically significant difference found between postoperative hypothermia and sex, age, preoperative temperature, magnitude of surgery; duration of anesthesia; duration of surgery, Operation Room temperature, Body Mass Index, fluid administration and blood transfusion.

## Discussion

Hypothermia is the most common complications in the recovery room which is associated with anesthesia and surgical intervention [15,32]. Early articles listed the incidence of hypothermia as ranging from 50% to 90% [33]. The current study showed that 30.72% patients had body temperature of less than 36°C on admission to the recovery room. This finding is lower than study conducted by Karalapillai [34], Kongsayreepong, et al. [19], Slotman et al. [35] and Abelha [26] which was 48.1%, 57.1%, 51% and 57.8% respectively. But it was high compared with study conducted by Shari [36] which is 4 percent. The result was almost comparable (32%) with study conducted by Clara Luis [37]. These differences could be due to small sample size, case mix, definition, site of measurement and thermometer type used as compared to the reference.

Inadvertent hypothermia is a recognized side effect of general anesthesia [38] because the body's normal thermoregulatory mechanisms, in particular vasoconstriction and shivering, are inhibited. While the mechanisms may differ, hypothermia also is an unintended side effect of regional anesthesia (i.e., spinal or epidural) [39]. Most patients in the present study underwent general anesthesia (68.5%) with was no correlation found between duration of anesthesia and postoperative temperature. This is in contrary with evidences, i.e. the longer the anesthesia, the lower the patient's body temperature [21]. However a statistically significant difference was observed between types of anesthesia and postoperative temperature. Accordingly, those who took general anesthesia had lower temperatures than spinal anesthesia (AOR, 1.84; 95% CI, 1.17; 3.24). This finding corroborates with literature, indicating that general anesthesia poses higher risk of postoperative hypothermia [20,26,40,41]. Regardless of the type of anesthesia selected, derangement of pre and intra- operative body temperature may have an impact on postoperative temperature. In this case, monitoring and maintaining patient body temperature are paramount [42].

Evidence also showed that the higher the American Society of Anesthesiology (ASA) physical status, the greater the risk for hypothermia [20,26,43,44]. Similarly this study showed higher ASA physical status had a significant risk factor for the development of postoperative hypothermia (AOR, 1.99; 95% CI, 1.16; 3.44).

Urgency of the procedure, especially emergency surgery, was also important risk factors for the development of postoperative hypothermia [20,26] which was in contrast with the current study finding. This showed that hypothermia should be considered in elective surgery too.

One of the limitations of the present study was the use of tympanic membrane temperature as a measurement of core temperature rather than the rectal one in which ear thermometer reading varies from 1 to 2 degrees below and above the temperature calculated by rectal and oral thermometers. Furthermore, this study includes different case mixes and procedures for body temperature control which may confound the results of this study. The other major limitation of this study was lack of generalizability.

## Conclusions and Recommendations

The incidence of postoperative hypothermia in recovery room was high. Patient with high ASA physical status, operated under general anesthesia and elective surgical patients were strongly associated with the development of postoperative hypothermia. Maintaining normothermia in these risky patients is important on patient safety as this reduces risk of complications, costs to the hospital and patients. All perioperative surgical team should participate to reduce the occurrence of postoperative hypothermia with early prediction and diagnosis for better outcome of patient.

## Authors' Contribution

Tadesse Belayneh participated in the conception, design, acquisition of the data, statistical analysis, interpretation of the data, drafting of the paper, and critical revision of the manuscript. Abewaw Gebeyehu and Zewditu Abdissa participated in the design and critical revision of the manuscript. All authors read and approved the final manuscript.

## Acknowledgment

The authors would like to thank University of Gondar (financial support), data collectors, and study participants.

## References

- Kempainen R, Brunette DD (2004) The evaluation and management of accidental hypothermia. *Respir Care* 49: 192-205.
- Burns SM, Wojnakowski M, Piotrowski K, Caraffa G (2009) Unintentional hypothermia: implications for perianesthesia nurses. *J Perianesth Nurs* 24: 167-173.

3. The management of inadvertent perioperative hypothermia in adults (2008) National Collaborating Centre for Nursing and Supportive Care commissioned by National Institute for Health and Clinical Excellence.
4. Sessler DI (2000) Perioperative heat balance. *Anesthesiology* 92: 578-596.
5. Snyder ML (2005) Learn the chilling facts about hypothermia. *Nursing* 35: 321-324.
6. Scott E, Buckland R (2006) A systematic review of intraoperative warming to prevent postoperative complications. *AORN J* 83: 1090-1104, 1107-13.
7. Good KK, Verble JA, Secrest J, Norwood BR (2006) Postoperative hypothermia-the chilling consequences. *AORN J* 83: 1054-1066.
8. Fried DJ (2004) Hypothermia's chilling effect on outcomes. *Nurs Manage* 35 Suppl OR Insider: 22-23.
9. Heier T, Caldwell JE (2006) Impact of hypothermia on the response to neuromuscular blocking drugs. *Anesthesiology* 104: 1070-1080.
10. Michelson AD, MacGregor H, Barnard MR, Kestin AS, Rohrer MJ, et al. (1994) Reversible inhibition of human platelet activation by hypothermia in vivo and in vitro. *Thromb Haemost* 71: 633-640.
11. Rajagopalan S, Mascha E, Na J, Sessler DI (2008) The effects of mild perioperative hypothermia on blood loss and transfusion requirement. *Anesthesiology* 108: 71-77.
12. Frank SM, Beattie C, Christopherson R, Norris EJ, Perler BA, et al. (1993) Unintentional hypothermia is associated with postoperative myocardial ischemia. The Perioperative Ischemia Randomized Anesthesia Trial Study Group. *Anesthesiology* 78: 468-476.
13. Panagiotis K, Maria P, Argiri P, Panagiotis S (2005) Is postanesthesia care unit length of stay increased in hypothermic patients? *AORN J* 81: 379-382, 385-92.
14. Mauermann WJ, Nemergut EC (2006) The anesthesiologist's role in the prevention of surgical site infections. *Anesthesiology* 105: 413-421.
15. Scott EM, Leaper DJ, Clark M, Kelly PJ (2001) Effects of warming therapy on pressure ulcers--a randomized trial. *AORN J* 73: 921-927, 929-33, 936-8.
16. Kurz A (2008) Thermal care in the perioperative period. *Best Pract Res Clin Anaesthesiol* 22: 39-62.
17. Kumar S, Wong PF, Melling AC, Leaper DJ (2005) Effects of perioperative hypothermia and warming in surgical practice. *Int Wound J* 2: 193-204.
18. Bilotta F, Pietropaoli P, La Rosa I, Spinelli F, Rosa G (2001) Effects of shivering prevention on haemodynamic and metabolic demands in hypothermic postoperative neurosurgical patients. *Anaesthesia* 56: 514-519.
19. Kongsayreepong S, Chaibundit C, Chadpaibool J, Komoltri C, Suraseranivongse S, et al. (2003) Predictor of core hypothermia and the surgical intensive care unit. *Anesth Analg* 96: 826-833, table of contents.
20. El-Rahmany HK, Frank SM, Vannier CA, Schneider G, Okasha AS, et al. (2000) Determinants of core temperature at the time of admission to intensive care following cardiac surgery. *J Clin Anesth* 12: 177-183.
21. Leslie K, Sessler DI (2003) Perioperative hypothermia in the high-risk surgical patient. *Best Pract Res Clin Anaesthesiol* 17: 485-498.
22. Kitamura A, Hoshino T, Kon T, Ogawa R (2000) Patient with diabetic neuropathy is at risk of a greater intraoperative reduction in core temperature. *Anesthesiology* 92: 1311-1318.
23. Dubois EF (1951) The many different temperatures of the human body and its parts. *West J Surg Obstet Gynecol* 59: 476-490.
24. Ikeda T, Sessler DI, Marder D, Xiong J (1997) Influence of thermoregulatory vasomotion and ambient temperature variation on the accuracy of core-temperature estimates by cutaneous liquid-crystal thermometers. *Anesthesiology* 86: 603-612.
25. Cattaneo CG, Frank SM, Hesel TW, El-Rahmany HK, Kim LJ, et al. (2000) The accuracy and precision of body temperature monitoring methods during regional and general anesthesia. *Anesth Analg* 90: 938-945.
26. Abelha FJ, Castro MA, Neves AM, Landeiro NM, Santos CC (2005) Hypothermia in a surgical intensive care unit. *BMC Anesthesiol* 5: 7.
27. Kelly G (2006) Body temperature variability (Part 1): a review of the history of body temperature and its variability due to site selection, biological rhythms, fitness, and aging. *Altern Med Rev* 11: 278-293.
28. Paes BF (2010) Accuracy of tympanic and infrared skin thermometers in children. *Arch Dis Child*.
29. Walpoth BH, Galdikas J, Leupi F, Muehleemann W, Schlaepfer P, et al. (1994) Assessment of hypothermia with a new "tympanic" thermometer. *J Clin Monit* 10: 91-96.
30. Chue AL, Moore RL, Cavey A, Ashley EA, Stepniewska K, et al. (2012) Comparability of tympanic and oral mercury thermometers at high ambient temperatures. *BMC Res Notes* 5: 356.
31. Kurz A (2008) Physiology of thermoregulation. *Best Pract Res Clin Anaesthesiol* 22: 627-644.
32. Hooper VD (2009) SPAN's evidence-based clinical practice guideline for the promotion of perioperative normothermia. *J Perianesth Nurs* 24: 271-287.
33. Burger L, Fitzpatrick J (2009) Prevention of inadvertent perioperative hypothermia. *Br J Nurs* 18: 1114, 1116-1119.
34. Karalapillai D, Story D (2008) Hypothermia on arrival in the intensive care unit after surgery. *Crit Care Resusc* 10: 116-119.
35. Slotman GJ, Jed EH, Burchard KW (1985) Adverse effects of hypothermia in postoperative patients. *Am J Surg* 149: 495-501.
36. Burns SM, Piotrowski K, Caraffa G, Wojnakowski M (2010) Incidence of postoperative hypothermia and the relationship to clinical variables. *J Perianesth Nurs* 25: 286-289.
37. Clara Luís, Carlos Moreno, Acácio Silva, Rosália Páscoa, Fernando Abelha (2012) Inadvertent Postoperative Hypothermia at Post-Anesthesia Care Unit: Incidence, Predictors and Outcome. *Open Journal of Anesthesiology* 2: 205-213.
38. Berry JM, Garrett K, Clifton GL (1999) Postoperative hypothermia in a tertiary care hospital. *Anesthesiology* 91: A1177.
39. Hart SR, Bordes B, Hart J, Corsino D, Harmon D (2011) Unintended perioperative hypothermia. *Ochsner J* 11: 259-270.
40. Biazotto CB, Brudniewski M, Schmidt AP, Auler Júnior JO (2006) Perioperative hypothermia. *Rev Bras Anestesiol* 56: 89-106.
41. Norris EJ, Beattie C, Perler BA (2001) Double masked randomized trial comparing alternate combinations of intraoperative anesthesia and anesthesia and postoperative analgesia in abdominal aortic surgery. *Anesthesiology* 95: 1054-1067.
42. Cattaneo CG, Frank SM, Hesel TW, El-Rahmany HK, Kim LJ, et al. (2000) The accuracy and precision of body temperature monitoring methods during regional and general anesthesia. *Anesth Analg* 90: 938-945.
43. Frank SM, El-Rahmany HK, Cattaneo CG, Barnes RA (2000) Predictors of hypothermia during spinal anesthesia. *Anesthesiology* 92: 1330-1334.
44. Lau LL, Hung CT, Chan CK, Chow B, Chui PT, et al. (2001) Anaesthetic clinical indicators in public hospitals providing anaesthetic care in Hong Kong: prospective study. *Hong Kong Med J* 7: 251-260.

**Citation:** Belayneh T, Gebeyehu A, Abdissa Z (2014) Post-operative Hypothermia in Surgical Patients at University of Gondar Hospital, Ethiopia. J Anesth Clin Res 5: 461. doi:[10.4172/2155-6148.1000461](https://doi.org/10.4172/2155-6148.1000461)